



## Genotoxicity of Aphicide Plus<sup>®</sup>, Eco Fruit-fly Bait GF120<sup>®</sup> and Rosecare 3<sup>®</sup> in the Allium cepa assay

Asita Okorie Asita\*, Relebohile Rebecca Mohale and Sibusisiwe Magama

Department of Biology, National University of Lesotho, P.O. Roma 180, Maseru 100, Lesotho, Southern Africa.

**Abstract:** Pesticides use boosts agricultural yield by reducing crop losses. However, some pesticides are mutagens and while technical grade active ingredients may produce mixed results in cytotoxicity and mutagenicity tests, synergistic interaction of pesticides, their metabolites or impurities in pesticide formulations often produce cytotoxic and genotoxic effects. *A. cepa* seedlings were treated for 24 hours, with three concentrations (mg mL<sup>-1</sup>) (0.04, 0.08 & 0.16) each of Aphicide Plus<sup>®</sup> (AP) (imidacloprid, 20g L<sup>-1</sup>); Eco Fruit-fly Bait GF120<sup>®</sup> (EF) (Spinosad, 0.24 g L<sup>-1</sup>) & Rosecare 3<sup>®</sup> (RC3) (bifenthrin, 2.0 g L<sup>-1</sup> & myclobutanil, 7.5 g L<sup>-1</sup>) and the effects on the (P+M)/(A+T) Ratio, cytotoxicity and genotoxicity were examined. The 0.08 mg mL<sup>-1</sup> of AP, 0.04 mg mL<sup>-1</sup> of EF, and 0.08 and 0.16 mg mL<sup>-1</sup> of RC3 induced changes in the (P+M)/(A+T) ratio, ( $p > 0.05$ ). The pesticides' concentrations tested depressed the mitotic index, ( $P < 0.05$ ) and were genotoxic, ( $P < 0.05$ ).

**Key words:** Insecticides; Fungicides; imidacloprid; Spinosad; bifenthrin; myclobutanil; Chromosome aberration.

### Introduction

In the field and during storage, agricultural products are threatened by insects, rodents, birds and other pests (Hayma, 2003). It was suggested by Raja *et al.* (2001), that insect damage in stored grains and other durable commodities may amount to 10-40% in developing countries, where modern storage technologies have not been introduced.

The control of pests that compete for food and fiber will continue to be an issue as the world population increases. According to Robson (2019) there are 7.7 billion people on the planet today and this number will grow to 9.7 billion in 2050. The need for food and fiber will continue and the need to control pests, using many means including agricultural chemicals will increase (Robson, 2019).

In order to cut down losses in agriculture, pesticides have been integrated into modern agricultural process to control weeds, diseases and insect pests that can markedly reduce the amount of harvestable produce and thereby increase outputs and productivity (Aktar *et al.*, 2009). Pimentel *et al.*, (1993) argued that, on average, the economic benefits from pesticide use are about four times their direct cost

to the users. About one-third of the agricultural products are produced by using pesticides (Liu *et al.*, 2002). Without pesticide application the loss of fruits, vegetables and cereals from pest injury would reach 78%, 54% and 32% respectively (Cai, 2008).

Pesticides are different from other agricultural inputs in that they do not directly boost yields in the way that fertilizers do; instead, they reduce crop losses caused by pests (Jha and Regmi, 2009).

Though pesticides use reduces crop losses and so increases agricultural output, pesticides are known to produce a wide spectrum of adverse health and environmental effects. The same pesticides that are effective in controlling serious harmful insects like the rice weevil (*Sitophilus oryzae*) and the European corn borer (*Ostrinia nubilalis*) are also lethal to beneficial insects like the honey bee (*Apis mellifera*) and pollinators, including the honey bee, are responsible for seventy-five percent of the world's food production Robson (2019). According to a report of WHO and UNEP, worldwide there are more than 26 million

\*Corresponding Author:

Asita Okorie Asita;

DOI- <http://dx.doi.org/10.21746/aps.2021.10.12.8>

human pesticide poisonings with about 220,000 deaths per year (Richter, 2002).

In a review of the literature on the health effects of pesticides, Mansour (2004) concluded that there is strong scientific evidence that pesticides, as a whole, can induce severe effects to human health ranging from myelotoxicity to cytogenetic damage and carcinogenicity. Other health effects of pesticides include acute and persistent damage in the nervous system (Kamel *et al.*, 2007), lung and respiratory disorders (Hoppin *et al.*, 2008), alterations in the reproductive organs (Hileman, 1994) birth defects (Rojas *et al.*, 2000).

Most of the adverse effects of pesticides to health, according to Norppa (2004) are the result of the genetic damage induced by genotoxic agents in somatic as well as in germinal cells and any genetic activity of chemicals, it has been suggested, is most likely to result from cell division abnormalities (Parry *et al.*, 1999).

Pesticides residues are known to persist in soil (Subbarao, 1999), water (Medina *et al.*, 1999) and in fruits and vegetables (Girotti *et al.*, 2009) and represent a risk for human health. Genotoxicity and mutagenicity of pesticides for non-target organisms and their influence on ecosystems are of worldwide concern (Pimentel *et al.*, 1998).

According to Zhang *et al.*, (2011), worldwide, there are currently about 500 pesticides used widely and in large quantities. In China alone, more than 400 companies are manufacturers of over 300 varieties of original pesticides and 3,000 formulations or commercial names. It was estimated that globally there were about 5.6 billion pounds of pesticide used annually with some 25 million agricultural workers poisoned (Jeyaratnam, 1990). These pesticides are manufactured as insecticides, fungicides, herbicides, rodenticides and antimicrobials to protect crops. In the southern African sub-region, a number of commercial formulations of pesticides are being marketed by South African Companies, three of which are Aphicide Plus® (AP) Eco Fruitfly

Bait GF-120® (EF) and Rosecare 3® (RC3) all manufactured by Efekto.

Aphicide Plus® (AP) is a systemic suspension concentrate contact and stomach insecticide for use in the home garden to control sucking insects, as indicated on the label, on conifers, roses and other ornamentals. It contains imidacloprid (chloro-nicotinyl) a neonicotinoid insecticide as the active ingredient. Imidacloprid acts on the central nervous system of insects as an agonist of the nicotinic acetylcholine receptor (Chao & Casida, 1997) stimulating the neurons and causing a subsequent fatigue and interference in the transmission of nerve impulses (Zhang *et al.*, 2000; Schulz-Jander and Casida, 2002). It is considered as an insecticide of low toxicity to mammals and other vertebrates due to its poor penetration of the blood-brain barrier of these organisms (Sheets, 2001 a, b). The U.S. Environmental Protection Agency (U.S. EPA) has classified imidacloprid into Group E, no evidence of carcinogenicity, based on studies with rats and mice. (Gervais, *et al.*, 2010; Thyssen and Machemer, 1999; Fed. Regist, 2005). However, following metabolic activation *in vitro*, imidacloprid produced calf thymus DNA adducts (Shah *et al.*, 1997) and was mutagenic in TA98 and TA100 *Salmonella typhimurium* strains, with or without S9 metabolic activation (Karabay and Oguz, 2005). Imidacloprid also induced significant increases in the frequency of sister chromatid exchange and micronuclei formation in human peripheral blood lymphocytes (Feng *et al.*, 2005; Costa *et al.*, 2009), mice and rat bone-marrow cells (Karabay and Oguz, 2005; Demisia *et al.*, 2007), and *Vicia faba* roots (Zang *et al.*, 2000).

Eco Fruit fly Bait GF-120® (EF), is selective concentrate bait for control of fruit fly species infesting various fruit and cucurbits. Eco Fruit fly Bait GF-120 contains the active ingredient spinosad (Naturalyte). Spinosad (SPN) is a naturally-occurring insecticide (larvicide) derived from the fermentation of a naturally occurring soil actinomycetes, *Saccharopolyspora spinosa* (Mertz and Yao, 1990; Mendonça *et al.*, 2019). SPN acts on different receptors of in-

sects, causing toxicological effects after ingestion or cuticle absorption. The mechanism of action of SPN is mainly through the nicotinic acetylcholine receptors and in a secondary pathway through the  $\gamma$ -amino-butyric receptors (GABA). As a consequence, SPN leads to hyper excitation, paralysis and death of the insect in the larval phase (Mendonça *et al.*, 2019). According to the World Health Organisation, the pesticide SPN has low acute toxicity and presents a low toxicological risk potential (WHO, 2005). It is used to control a wide variety of pests. These include thrips, leaf miners, spider mites, mosquitoes, ants, fruit flies and others and has been registered for use in pesticides by the U.S. EPA since 1997 (Bunch *et al.*, 2014). Spinosad was not carcinogenic to Fischer 344 rats at dose levels up to 0.05% (Yano *et al.*, 2002). There are conflicting reports on the mutagenicity of spinosad.

Spinosad has been reported to lack a mutagenic effect by U.S. EPA (1997) while Mansour *et al.*, (2008) found that spinosad was genotoxic to rat bone marrow cells. Spinosad lacked genotoxic activity in the *in vivo* somatic mutation and recombination test (SMART) in *D. melanogaster* (Akmoutsou *et al.*, 2011). According to Marrs (2012), none of the genotoxicity studies showed mutagenic activity associated with spinosad. Chronic exposure studies failed to induce tumor formation in rats and mice; mice given up to 51 mg/kg/day for 18 months resulted in no tumor formation (Stebbins, 2002).

Rosecare 3<sup>®</sup> (RC3) is a contact insecticide plus a systemic fungicide formulated as a micro emulsion for the control of diseases including aphids, red spider mite, thrips, black spot, rust and powdery mildew on roses and other ornamentals. (<https://www.hadeco.co.za/>). Rosecare 3<sup>®</sup> contains bifenthrin and myclobutanil as the active ingredients. Bifenthrin is a synthetic pyrethroid insecticide in the pyrethroid family constituting man-made versions of pyrethrins, produced naturally by chrysanthemum flowers (Johnson *et al.*, 2010) and myclobutanil is a triazole fungicide. The insecticide bifenthrin is designed to be effective

by contact or ingestion and affects the central and peripheral nervous systems of invertebrates thereby killing them (Tomlin, 2000). The U.S. EPA classified bifenthrin as a Category C, possible human carcinogen, based on studies in mice. Other studies indicate that bifenthrin does not cause cancer or birth defects when fed to rats or rabbits that ate bifenthrin when pregnant (Johnson *et al.*, 2010). A "FAO/WHO Joint Meeting on Pesticide Residues" (JMPR) examined the test battery for the assessment of mutagenicity of bifenthrin and concluded that the results were mixed (WHO, 2017). The JMPR concluded that, whereas some tests were clearly negative (such as the Ames test on different strains of *Salmonella typhimurium* with and without metabolic activation), other tests, such as the Mouse Lymphoma Mutagenesis Assay or the unscheduled DNA synthesis test with rat hepatocytes showed weak positive response or yielded inconclusive results with technical bifenthrin (WHO, 2017).

Myclobutanil, according to the European Food Safety Authority (EFSA) is a systemic fungicide with preventive, curative and eradicator properties that belongs to the class of conazole fungicides approved for use in the EU and many other countries (EFSA, 2010). Conazoles are a diverse group of commercially important fungicides with clinical and agricultural applications for cereals' treatment, vegetables, fruits and flowers and also as medical products (Ross *et al.*, 2009; Hester *et al.*, 2012). Myclobutanil works by inhibiting the biosynthesis of ergosterol, a critical component of fungal cell membranes. If these membranes are unable to grow, the growth of the fungus is inhibited as well ([domyown.com/myclobutinil-c-114\\_468.html](http://domyown.com/myclobutinil-c-114_468.html)). Myclobutanil is moderately toxic to birds, fish, aquatic invertebrates, algae, honeybees and earthworms (Lewis *et al.*, 2016) It has been shown that the toxicity of triazole fungicides passes through oxidative stress, causing DNA damage or apoptosis (Ross *et al.*, 2012; Hester *et al.*, 2006). However, Myclobutanil did not show any genotoxic potential in both *in vitro* and *in vivo* genotoxicity tests (EFSA, 2010).

Examination of the literature on the genotoxicity and mutagenicity of the technical grade active ingredients (TGAIs) of the three pesticides (imidacloprid, spinosad, bifenthrin, and myclobutanil) reveal conflicting reports, but we could not find reports of tests using the *Allium cepa* chromosome aberration assay for some. The present study aimed therefore to investigate the cytotoxicity and genotoxicity of different concentrations of three pesticide formulations (Aphicide Plus®, Eco Fruit-fly Bait GF-120® and Rosecare 3®), containing these active ingredients using the *Allium cepa*, chromosome aberration assay because synergistic interaction or potentiation between or among pesticides that are mixed together is a well-known phenomenon (Cloyd *et al.*, 2007).

The *Allium cepa* L assay is one of the established plant bioassays, validated by the international programme on chemical safety (IPCS, WHO), as an efficient and standard test for chemicals screening, *in situ* monitoring of the genotoxicity of environmental substances (Leme and Marin-Morales, 2009). The *Allium cepa* L assay is an *in vivo* assay that tests genotoxicity using chromosomes and therefore detects chromosome structural and numerical alterations (Tedesco and Laughinghouse, 2012; Bonciu *et al.*, 2018). The results obtained, using the *Allium cepa* assay have been shown to be similar with those of mammalian and non-mammalian test systems (Constantin and Owen, 1982; Fiskesjö, 1985; Cauhan *et al.*, 1999; Aydemir *et al.*, 2008). *Allium cepa* root cells, just like many plant cells, have the monooxygenase enzyme systems (MFO-system) and therefore the ability to activate promutagens (Plewa and Gentile, 1982; Higashi, 1988; Fiskesjö, 1985).

## Material and Methods

### Test organism

Seeds of *Allium cepa* variety, Texas Grano 502 P.R.R®, obtained from Sakata seeds, Lanseria 1748, South Africa were used for the study.

### Pesticides

Pesticides used in the study were products of Efekto; Aphicide Plus® Reg. No. L8780 Act

No. 36 of 1947 N-AR1162 (contained imidacloprid at 20 g L<sup>-1</sup> in the formulation; Manufacturer's recommendation, 10 - 20 mL L<sup>-1</sup> of water = 101 - 51 x dilution); Eco Fruitfly Bait GF-120® model number 6001379103341 (contained spinosad (naturalyte) at 0.24 g L<sup>-1</sup>; Manufacturer's recommendation, 1.2 ml in 9 to 30L of water = 8.5 - 26 x dilution); Rosecare 3® registration number L7599 N-AR 0726 Act No. 36 of 1947 (contained bifenthrin at 2.0 g L<sup>-1</sup> and myclobutanil at 7.5 g L<sup>-1</sup> in the formulation; Manufacturer's recommendation, 10ml per 1L of water = 101 x dilution). All the pesticides were products of Agro-Serve (Pty) Ltd, South Africa. The pesticides were purchased from the Maseru Garden Centre, Lesotho, Southern Africa.

### Chemicals

Methanol (Absolute) was a product of Associated Chemical Enterprises (PTY) LTD of The Republic of South Africa; Hydrochloric acid and Glacial acetic acid were products of UNILAB of The Republic of South Africa; Aceto-carmin stain from Carolina Biological Supply Company, USA.

### Genotoxicity Experiments

The preliminary assay to select concentrations of pesticides to use for the Genotoxicity (GT) assay (including the treatment of *Allium cepa* seedlings with pesticides, root harvest, slide preparation and scoring of slides) were conducted according to the methods of Asita *et al.* (2017). From the results of the preliminary assays to select the concentrations of pesticides to use, the following concentrations of pesticides (in mgmL<sup>-1</sup>) AP (0.04, 0.08 and 0.16); EF (0.04, 0.08 and 0.16); and RC3 (0.04, 0.08 and 0.16) were assessed for cytotoxicity that is, mitotic index (MI) and genotoxicity (GT) following 24 hours of treatment of the *A. cepa* seedlings. When diluted in accordance with the manufacturers' recommendation, the concentrations of the individual active ingredient in the applied pesticide will be: imidacloprid at 20 g L<sup>-1</sup> in formulation and at 101 - 51 x dilution = 20/101 - 20/51 = 0.198 - 0.392 mg mL<sup>-1</sup> Spinosad at 0.24 g L<sup>-1</sup> in the formulation and at 8.5 - 26 x dilution =

$0.24/8.5 - 0.24/26 = 0.028 - 0.00923 \text{ mg mL}^{-1}$ ; Bifenthrin at  $2 \text{ g L}^{-1}$  in the formulation and at  $101 \times$  dilution =  $2/101 = 0.02 \text{ mg mL}^{-1}$ ; myclobutanil at  $7.5 \text{ g L}^{-1}$  in the formulation and at  $101 \times$  dilution =  $7.5/101 = 0.074 \text{ mg mL}^{-1}$ . The tested concentrations (0.04, 0.08 and 0.16) are lower than the applied concentration of imidacloprid, a bit higher than the recommended concentration for use for spinosad and bifenthrin but similar to the recommended concentration for use for myclobutanil. In each assay, three root tips (triplicate) were assessed at each concentration. On each of three slides ( $n = 3$ ) per treatment, a total of 2000 cells, classified into interphase or dividing cell, that is, prophase (normal, N or aberrant, ABN), metaphase (N or ABN), anaphase (N or ABN) or telophase (N or ABN) were scored. The aberrant (ABN) category in each cell division stage comprised of cells having their mitotic division apparatus (Chromosomes, spindle fibres, kinetochore or centrosomes) damaged. Therefore, a total of 6000 cells each were examined and scored for the negative control (water) and treatment (pesticide concentration) groups.

Mitotic stage cells containing the following types of damages were classified as aberrant (ABN) (Asita *et al.*, 2017):

Chromosome fragments (F) - piece of chromosome broken from whole chromosome as a result of pesticide treatment and lacking centromere.

Anaphase or Telophase bridge (A.B) - Dicentric chromosomes that form a bridge between both poles at anaphase or telophase. Often it indicates paracentric inversions or other possibilities that include breakage and fusion of chromosomes and sister chromatid reunion.

Laggard (L) - whole chromosomes that fail to migrate to either pole at anaphase because of damage to the kinetochore,

C-Mitosis (C-Mit) - Mitotic cells that lack spindle fibres so that the chromosomes lie scattered throughout the cell. The effect is usually produced in cells treated with the

spindle poisons, colchicines or colcemid, hence C-Mitosis.

Sticky chromosomes (S) - sticky chromosomes fail to condense completely so that at metaphase, the chromosomes are still long like prophase chromosomes and remain entangled with each other. In extreme cases, a chromatin mass, undistinguishable as chromosomes is seen as a clump. If such damage occurs in interphase cells, they are referred to as pyknotic cells.

### Analysis of Slide Preparations

#### Cytotoxicity

The mitotic index (MI) was expressed as the number of dividing cells per 100 cells scored according to the formula:

$$\text{MI} = \text{Number of dividing cells} / \text{Total number of cells scored} \times 100. \quad (2)$$

The MI was used as a measure of cytotoxicity (CT). The MI of each treatment group was compared with that of the negative control group using t-test at a probability level of 0.05, using the SPSS for windows, version 11.0 software.

#### Genotoxicity

Genotoxicity (GT) was expressed as the number of aberrant mitotic cells (AMC) per 100 mitotic cells [*i.e.* AMC + normal mitotic cells (NMC)] scored according to the formula:

$$\text{Frequency of GT} = \text{AMC} / (\text{AMC} + \text{NMC}) \times 100 \quad (3)$$

The mean GT of each group of three slides per concentration of test agent was compared with that of the negative control group using t-test. P values less than 0.05 ( $P < 0.05$ ) were considered as indicative of significance.

#### Data Analysis


Data were expressed as mean  $\pm$  SD of three values. For the determination of cytotoxicity and genotoxicity, the mean value of each group of three slides per concentration of test pesticide, was compared with that of the negative control group using student's t-test. P-values less than 0.05 ( $p < 0.05$ ) were considered as indicative of significance. Analysis

was performed using the SPSS for windows version 11.0 software.

## Results

### Cytotoxicity (CT) & Genotoxicity (GT) of the Pesticides in the *A. Cepa* Root Tip Chromosome Aberration Assay

Photographs of the most representative pictures of normal mitotic cells and cells containing the different types of chromosome aberrations that were observed and scored are presented in Figure 1.



**Fig. 1.** Photomicrographs of cells of *Allium cepa* showing untreated cells in normal division stages (First row) and Chromosomal aberrations (arrowed) in cells treated with pesticides. (a) Interphase (b) Normal Prophase (c) Normal metaphase (d) Normal metaphase (e) Early anaphase (f) Late anaphase (g) Telophase (h) Pyknotic interphase nuclei with micronucleus (i) Prophase with sticky chromosomes (j) Metaphase with sticky chromosomes (k) Metaphase with sticky chromosomes (l) C-metaphase (m) Metaphase with dislocated chromosome (n) Late anaphase with dislocated chromosome (o) Anaphase with sticky and scattered chromosomes (p) Late anaphase with chromosome bridge (q) Telophase with lagging chromosome (r) Telo-

phase with sticky chromosomes and bridge (s) telophase with chromosome bridge and lagging (t) telophase with chromosome bridge and fragment (u) Telophase with chromosome fragment and lagging. Magnification is 1000 X.

### Cytotoxicity and Genotoxicity of the Pesticides

The results of the cytotoxicity (effects on the mitotic index) and genotoxicity (damage to mitotic apparatus) of the treatment of onion root tip meristem cells with different concentrations of the pesticides are presented in Table 1.

**Table 1.** Toxic and Genotoxic effects of Aphicide Plus®, Eco Fruit-fly Bait GF-120® and Rosecare 3® on onion root tip meristem cells following 24h exposure to three different concentrations of each pesticide

c	Conc. (mg/ml)	Statistics	Interph	Cells in division stages per 2000 cells scored										Total Cells Scored	Prophase + Metaphase and Anaphase + Telophase Analysis			MI	GENOTOXICITY	
				Proph		Metaph		Anaph		Teloph		Total			P + M	A + T	(P+M)/(A+T)			
				N	A	N	A	N	A	N	A	N	A							N
Water	Water	Mean	1790.00	1	1	2	1	6	0	5	0	2	2	2	20	15	5	2.6	10	1.05
		S.D	75.54	6	1	8	1	1	0	2	0	7	0	7	0	55	2	0.1	3	0.41
Aphicide Plus	0.04	Mean	1920.33	3	4	1	2	3	0	1	1	7	8	8	20	59	2	2.9	3	10.38*
		S.D	13.65	2	2	2	2	6	0	1	1	1	2	1	0	24	4	0.5	0	3.55
	0.08	Mean	1978.67	2	1	1	2	0	0	1	3	4	1	2	20	17	3	8.1*	1	86.90*
		S.D	5.86	4	3	1	1	0	0	1	2	6	1	6	0	2	3	6.9	0	22.68
0.16	Mean	1962.00	1	8	6	3	1	0	6	2	2	1	3	20	29	9	3.3	2	33.35*	
	S.D	14.00	1	1	1	1	1	0	0	1	2	1	2	0	0	2	0.8	0	0.86	
Eco Fruit-fly Bait GF120	0.04	Mean	1900.67	5	1	8	2	1	1	1	8	7	2	9	20	78	2	3.8*	5	23.49*
		S.D	7.37	2	4	2	1	1	1	4	3	6	3	7	0	5	2	0.2	0	2.63
	0.08	Mean	1972.33	9	7	3	2	0	0	6	4	1	1	2	20	21	9	2.5	1	48.62*
		S.D	6.51	2	3	1	1	0	1	2	3	4	3	7	0	5	5	0.6	0	3.27

Rosecare3	0.16	Me an	19 86. 67	2	6	2	0	0	0	2	1	6	8	1 3	20 00	10 .3	3. 0	4.6	1. 00 *	57.90*
		S.D	2.5 2	2	3	1	1	0	1	1	1	3	3	3	0	4. 5	1. 7	3.2	0. 00	17.59
	0.04	Me an	19 26. 33	3 6	7	1 3	3	0	1	1 2	2	6 0	1 3	7 4	20 00	59 .0	1 4. 7	4.6	3. 67 *	18.01*
		S.D	13. 87	8	2	4	2	0	1	6 1	1	5 2	1 5	1 4	0	7. 0	7. 0	2.0	0. 58	5.78
	0.08	Me an	19 44. 33	2 0	1 0	5	2	0	1	1 1	7	3 6	2 0	5 6	20 00	37 .0	1 8. 7	2.0*	2. 67 *	35.21*
		S.D	11. 93	6	3	2	1	0	1	3 2	9	6	1 6	1 2	0	7. 2	4. 9	0.2	0. 57	6.67
	0.16	Me an	19 59. 67	1 0	1 0	2	3	0	0	5 1	1 0	1 7	2 3	4 0	20 00	25 .3	1 5. 0	1.7*	2. 00 *	57.64*
		S.D	5.5 1	5	5	1	4	0	0	3 1	1	5 6	6	6	0	6. 1	2. 0	0.3	0. 00	11.09

TC = Test Compound; Amphicide Plus®; Eco Fruit-fly Bait, GF120®; Rosecare 3®; MI = Mitotic Index; Interph = Interphase; Proph = Prophase; Metaph = Metaphase; Anaph = Anaphase; Teloph = Telophase; A&T = (Anaphase & Telophase); S.D = Standard deviation; \* Significant difference from control in the t-test at P<0.05 and 4 d.f.

### The (P+M)/ (A+T) Ratio, Cytotoxicity and Genotoxicity Induced by the Pesticides

The following results were observed following the treatment of root tips of *A. cepa* seedlings with Aphicide Plus® (0.16 mg mL<sup>-1</sup>, 0.08 mg mL<sup>-1</sup> and 0.04 mg mL<sup>-1</sup>), Eco Fruit-fly GF120® (0.16 mg mL<sup>-1</sup>, 0.08 mg mL<sup>-1</sup> and 0.04 mg mL<sup>-1</sup>), Rosecare 3® (0.16 mg mL<sup>-1</sup>, 0.08 mg mL<sup>-1</sup> and 0.04 mg mL<sup>-1</sup>) for 24 hours.

**(P+M)/ (A+T) Ratio:** Examination of the (P+M)/(A+T) ratio in column 19 of Table 1 shows that only the middle concentration (0.08 mg mL<sup>-1</sup>) of Aphicide Plus®, the lowest concentration (0.04 mg mL<sup>-1</sup>) of Eco Fruit-fly Bait, GF120® and the middle and highest concentrations (0.08 and 0.16 mg mL<sup>-1</sup>) of Rosecare® 3 induced significant change in (P+M)/(A+T) ratio, when compared with the water treated negative control group ( $p > 0.05$ ).

**Cytotoxicity:** Examination of the MI in column 20 of Table 1 shows that all three concentrations of each of the three pesticides tested induced a significant reduction of the MI when compared to the solvent (water) treated negative control ( $P < 0.05$ ) and were adjudged toxic to the root meristem cells of *A. cepa*.

**Genotoxicity (GT):** Examination of induction of genotoxicity in column 21 of Table 4 shows that all three concentrations of each of the three pesticides tested induced a significant increase in genotoxicity when compared to the solvent (water) treated negative control ( $P < 0.05$ ) and were adjudged genotoxic to the root meristem cells of *A. cepa*.

### Discussion

In this study, three concentrations (mg mL<sup>-1</sup>) each of three pesticides, Aphicide Plus® (0.16, 0.08 and 0.04), Eco Fruit-fly GF120® (0.16, 0.08 and 0.04), Rosecare 3® (0.16, 0.08 and 0.04) were assessed for their effect on the (P+M)/(A+T) Ratio, cytotoxicity and genotoxicity using the onion (*Allium cepa* L.) root tip meristem chromosome aberration assay system. The results of the effects of the pesticides on

the three endpoints following the treatment of onion root tip meristem cells with different concentrations of the pesticides are presented in Table 1.

The (P+M)/(A+T) Ratio, cytotoxicity and genotoxicity results following treatment of *Allium cepa* L root tip meristem cells with Aphicide Plus® (imidacloprid (chloro-nicotinyl) at 20g L<sup>-1</sup> in the formulation) are presented in Table 1, rows 6 to 11. Only the middle concentration (0.08 mg/mL) of Aphicide Plus®, induced significant increase in the relative proportions (P+M) to (A+T) ratio, when compared with the water treated negative control group ( $p > 0.05$ ). The lowest and highest concentrations (0.04 and 0.16 mg mL<sup>-1</sup> respectively) increased the (P+M)/(A+T) ratio also but the increases were not significant. A decrease in the proportion of dividing cells in A+T is an indication of metaphase arrest due to the poisoning of the spindle fibres, akin to the action of the well documented spindle poison, colcemid (Parry *et al.*, 1999). The chemotherapeutic agents taxol, vincristine, vinblastine and nocodazole act similarly (Alberts *et al.*, 2008). These compounds act by binding to and stabilizing microtubules, inhibiting their dynamic instability and causing various genetic disruptions, including the induction of cell cycle arrest (Alberts *et al.*, 2008; Zhang *et al.*, 2015). Since only one concentration modified the (P+M)/(A+T) ratio, it is concluded that Aphicide Plus® could be a spindle fibres poison. All three concentrations of Aphicide Plus® tested reduced the mitotic index (MI) of treated root tip cell populations. Mitotic index is considered a parameter that allows one to estimate the frequency of cellular division (Marcano *et al.*, 2004) and the reduction of mitotic activities has been used frequently to trace substances that are cytotoxic (Linnainmaa *et al.*, 1978; Smaka-Kincl *et al.*, 1996). Many investigators have recorded a depression of the mitotic index following the treatment of test organisms with pesticides (Panda and Sahu, 1985; Amer and Farah, 1974). In the

present study, treatment of the *A. cepa* root tips with each of the three concentrations of Aphicide Plus® tested reduced the MI which indicated a reduction of cell division or proliferation, compared to the water treated negative control. It is concluded that the concentrations of Aphicide Plus® tested in the present study are cytotoxic since they modify normal MI, in this case, decreased the MI (Leme and Marin-Morales, 2009; Vieira and Silveira, 2018). Aqueous extracts of leaves and stems of the plant *P. barbatus*, increased the cell division of *A. cepa* roots (Iganici et al., 2006; Trapp et al., 2020). Similarly, all three concentrations of Aphicide Plus® tested induced genotoxicity. In studies performed by Karabay and Oguz (2005), they observed significant levels of chromosome aberration (CA) (breaks and chromosomal adherences) in erythrocytes of mice exposed to imidacloprid, the active ingredient in Aphicide Plus®, and also in the studies of Rodríguez et al. (2015), in which they observed genotoxicity for imidacloprid which induced different kinds of CA in *A. cepa* cells, including binucleated cells, losses, breaks, chromosomal bridges, and MN. Imidacloprid (0.036, 0.36 & 3.6 g L<sup>-1</sup>) was also cytotoxic, reducing the MI and genotoxic in the *A. cepa* chromosome aberrations assay as it induced different types of CA, mainly bridges and chromosomal adherences (Bianchi et al., 2016). However, some studies which investigated the genotoxic effects of imidacloprid at several concentrations and different organisms have produced conflicting results, both positive and negative (Solecki, 2001; Feng et al., 2004; Karabay and Oguz, 2005; Demisia et al., 2007; Jemec et al., 2007; Kreutzweiser et al., 2007; Rodríguez et al., 2015). Synergistic interaction or potentiation between or among pesticides that are mixed together is a well-known phenomenon (Cloyd et al., 2007). In a recent study (Ilyushina et al., 2020), using technical grade active ingredients (TGAIs), imidacloprid, imazalil and tebuconazole pesticides individually at doses of up to 120, 300 and 1000 mg kg<sup>-1</sup> bwd-

<sup>1</sup>, respectively, did not induce micronucleus formation in PCE of mice bone marrow in CD-1 mice, whereas their combination in a ratio of 14.0/1.7/1.0 by weight demonstrated negative results in the Ames (*Salmonella typhimurium*) test but induced a statistically significant, dose-dependent increase in micronuclei (MN) in polychromatic erythrocytes (PCEs) in mouse bone marrow at doses lower than those used separately but without suppressing erythropoiesis, meaning that it was not cytotoxic. They concluded that the observed effect may be mediated by the synergistic action of the tested pesticides, their metabolites or impurities. The synergism of endosulfan and chlorpyrifos was shown in cultured human peripheral blood lymphocytes by chromosomal aberration test and comet assay (Sultana Shaik et al., 2016). In the present study, the concentration of imidacloprid of 20g L<sup>-1</sup> or 20 mg mL<sup>-1</sup> in Aphicide Plus® formulation is much higher than the tested concentrations (0.16, 0.08 & 0.04 mg mL<sup>-1</sup>). These very low concentrations still induced cytotoxicity and genotoxicity. The observed cytotoxic and genotoxic effects of the low concentrations of Aphicide Plus® may thus be attributed to the synergistic action of components of the Aphicide Plus®, their metabolites or impurities. The genotoxic effect of Aphicide Plus® observed in the present study demonstrated the genotoxic effect of imidacloprid containing pesticide formulation.

The (P+M)/(A+T) Ratio, cytotoxicity and genotoxicity results following treatment of *Allium cepa* L root tip meristem cells with Eco Fruit-fly Bait GF-120® (Spinosad (Naturalyte) at 0.24 g L<sup>-1</sup> in the formulation) are presented in Table 1, rows 12 to 17. Only the lowest concentration (0.04 mg mL<sup>-1</sup>) of Eco Fruit-fly Bait GF120®, induced significant increase in the relative proportions P+M and A+T ratio, when compared with the water treated negative control group (p > 0.05). All the three concentrations (0.04, 0.08 and 0.16 mg mL<sup>-1</sup>) of

Eco Fruit-fly Bait GF-120® tested were cytotoxic and genotoxic.

There are conflicting reports on the genotoxicity/mutagenicity of spinosad in earlier studies. Spinosad was reported to lack a mutagenic effect by U.S. EPA (1997) while Mansour et al. (2008) found that spinosad was genotoxic to rat bone marrow cells, but was not genotoxic in the *in vivo* somatic mutation and recombination test (SMART) in *D. melanogaster* (Akmoutsou, et al., 2011). In a study that investigated the genotoxicity of dillapiol and spinosad (0.31, 0.96 & 1.6 µg mL<sup>-1</sup>) in distilled water as solvent, using the somatic mutation and recombination test (SMART) in wings of *Drosophila melanogaster*, both compounds were genotoxic, inducing mainly mitotic recombination events (Aciole et al., 2014). In a recent study, spinosad (SPN) at the concentrations of 0.625; 1.25 and 2.5 g L<sup>-1</sup> induced a high frequency of micronuclei in *Tradescantia pallida* (*T. pallida*) and was adjudged genotoxic, whereas, in the same study, 0.039; 0.078 and 0.156 µg mL<sup>-1</sup> of spinosad were not mutagenic in the *in vivo* somatic mutation and recombination test (SMART) in *D. melanogaster* (Mendonça et al., 2019). The assessment of pure spinosad for genotoxicity/mutagenicity using different test systems, *in vitro* and *in vivo*, as shown in the studies reported above, presents conflicting results. In the present study which used the *A. cepa in vivo* chromosome aberration test to evaluate Eco Fruit-fly Bait GF-120® (spinosad at 0.24 g L<sup>-1</sup> or 0.24 mg mL<sup>-1</sup> in the formulation) for cytotoxicity and genotoxicity, all the three concentrations (0.04, 0.08 & 0.16 mg mL<sup>-1</sup>) tested were cytotoxic and genotoxic. The unequivocal genotoxic effect of Eco Fruit-fly Bait GF-120® observed in the present study demonstrates the genotoxic effect of spinosad containing pesticide formulation and suggests a possible effect of synergistic interaction of the contents of the pesticide formulation.

The (P+M)/(A+T) Ratio, cytotoxicity and genotoxicity results following treatment of *Allium cepa* L root tip meristem cells with Rosecare 3®

(bifenthrin at 2.0 g L<sup>-1</sup> & myclobutanil (a triazole) at 7.5 g L<sup>-1</sup> in the formulation) are presented in Table 1, rows 18 to 23. The middle and highest concentration (0.08 and 0.16 mg mL<sup>-1</sup>) of Rosecare 3®, induced significant increase in the relative proportions (P+M) to (A+T) ratio, when compared with the water treated negative control group (p > 0.05). All the three concentrations (0.04, 0.08 & 0.16 mg mL<sup>-1</sup>) of Rosecare 3® tested were cytotoxic and genotoxic. Reports of mutagenicity and genotoxicity studies of bifenthrin using different assay test systems are mixed, some positive while others are negative. Bifenthrin was not mutagenic in mutagenicity assays including the Ames test, *in vivo* rat bone marrow cells, Chinese hamster ovary (CHO) cells, and unscheduled DNA synthesis (UDS) at concentrations of bifenthrin up to 2.5 µl mL<sup>-1</sup>. However, bifenthrin showed mutagenic properties in a mouse lymphoma L5178 Y cells assay (U.S. Environmental Protection Agency, 1988; Johnson et al., 2010). Bifenthrin (technical grade-purity 95%) concentrations (0.016, 0.0347 and 0.0628 ppm) caused different types of chromosome aberrations in livers cells of the fish, *Channa punctatus* (Chaudhari and Saxena, 2015) and DNA damage in gill tissue of the fresh water fish, *Danio rerio* when assessed using the alkaline comet assay (Reddy et al., 2013). The Rosecare 3® evaluated in the present study is a formulation containing bifenthrin at 2.0 g L<sup>-1</sup> (2 mg mL<sup>-1</sup>) and was unequivocally cytotoxic and genotoxic in the *A. cepa* chromosome aberration assay. However, a 225 mg L<sup>-1</sup> mixture of dichloran (0.24); dichlofluanid (0.03); iprodione (2.97); chlorothalonil (0.08) and bifenthrin (1.10) caused significant increase in the frequency of micronuclei, reduced mitotic index and increased anaphase chromosome aberrations in *Allium cepa* root (Feretti et al., 2007). Reports of mutagenicity and genotoxicity studies of myclobutanil using different assay test systems are mixed, some positive while others are negative. Myclobutanil was not genotoxic in the *in vitro* Ames Bacterial reverse mutation assay, did

not induce chromosomal aberrations with and without metabolic activation up to 200  $\mu\text{g mL}^{-1}$  in the *in vitro* cytogenetics structural Chromosomal Aberration Assay, was not genotoxic in the CHO/HGPRT assay nor in the *in vivo* Mouse Micronucleus test and did not induce an increase in unscheduled DNA synthesis up to toxic dose. 0.1-1000  $\text{g mL}^{-1}$  tested (U.S. EPA, 2000; EFSA, 2010). Myclobutanil did not induce mutations in the Big Blue mouse liver *in vivo* assay system (Ross *et al.*, 2009) and was not cytotoxic nor genotoxic to four human cell lines (ACHN, SH-SY5Y, LS-174T, HepG2) when screened with the  $\gamma\text{H2AX}$  In Cell Western (ICW) assay that simultaneously determines cytotoxic and genotoxicity of xenobiotics on cells cultured in a 96-well plate format after 24 h of treatment (Graillot *et al.*, 2012). The Rosecare 3<sup>®</sup> evaluated in the present study is a formulation containing myclobutanil (7.5  $\text{g L}^{-1}$  or 7.5  $\text{mg mL}^{-1}$ ) and was unequivocally cytotoxic and genotoxic in the *A. cepa* chromosome aberration assay. A 225  $\text{mg L}^{-1}$  mixture of either, dimethoate (0.10); vinclozolin (0.92); procymidone (1.44); fenarimol (0.03); hexaconazole (0.12); myclobutanil (0.27) or parathion-ethyl (0.11); folpet (1.03); procymidone (0.14); acephate (0.05); myclobutanil (0.49) caused significant increase in the frequency of micronuclei, significant reduction in mitotic index and increase in anaphase chromosome aberrations in *Allium cepa* root (Feretti *et al.*, 2007). The unequivocal genotoxic effects of Rosecare 3<sup>®</sup> observed in the present study demonstrate the genotoxic effect of bifenthrin and myclobutanil containing pesticide formulation and suggest a possible effect of synergistic interaction of the contents of pesticide formulation.

Treatment with each of all the three pesticides caused a dose related cytotoxicity (reduction of the MI) without any real direct action on the spindle, with the exception of the treatment with 0.08  $\text{mg mL}^{-1}$ . Aphicide Plus<sup>®</sup>, 0.04  $\text{mg mL}^{-1}$  of Eco Fruit-fly Bait GF120<sup>®</sup> and 0.08 and 0.16  $\text{mg/mL}$  of Rosecare 3<sup>®</sup>. The mitotic index

decreased with increased dose and at the same time, the relative proportions of P+M and A+T types remained the same. This may be taken as a typical indicator of cytotoxic damage (Parry *et al.*, 1999).

The abnormal dividing or mitotic cells that were induced by the pesticides and scored in the present study were prophase, metaphase, anaphase and telophase cells with sticky chromosomes, c-metaphase, cells with dislocated chromosomes, cells with chromosome bridge and or lagging chromosome and cells with chromosome fragments.

Stickiness has been shown to be as a result of DNA condensation (Österberg *et al.*, 1984) and entanglement of inter-chromosomal chromatin fibers which led to subchromatid connections between chromosomes (Patil and Bhat, 1992). Levan (1938) described colchicine mitosis (c-metaphase or c-anaphase) as an inactivation of the spindle followed by a random scattering of the condensed chromosomes in the cell. According to Yildiz and Arikan (2008), a large number of laggard chromosomes and C-anaphases indicate a test compound acted as a potent spindle inhibitor. The induction of vagrant chromosomes according to Elghamery *et al.* (2003), leads to the separation of unequal number of chromosomes in the daughter nuclei and subsequently formation of daughter cells with unequal sized or irregularly shaped nuclei at interphase. The induction of anaphase/telophase bridges has been attributed to chromosome breaks, stickiness and breakage and reunion of the broken ends (Parry *et al.*, 1985; Badr *et al.*, 1992).

### Conclusion

Of the three concentrations ( $\text{mg mL}^{-1}$ ) (0.04, 0.08 and 0.16) of each pesticide tested, the 0.08  $\text{mg mL}^{-1}$  of Aphicide Plus<sup>®</sup>, 0.04  $\text{mg/mL}$  of Eco Fruit-fly Bait GF120<sup>®</sup> and 0.08 and 0.16  $\text{mg mL}^{-1}$  of Rosecare 3<sup>®</sup> induced significant change in (P+M)/(A+T) ratio, ( $p > 0.05$ ). All three concentrations of each pesticide significantly depressed the mitotic index (MI) and

were adjudged cytotoxic ( $P < 0.05$ ). Each concentration of the three pesticides tested, induced genotoxicity ( $P < 0.05$ ). The aberrations induced were, sticky chromosomes, c-metaphase, dislocated chromosomes, Chromosome bridges, lagging chromosomes and chromosome fragments. Whereas genotoxicity tests of some of the pure compounds yielded mixed results in different test systems, the three concentrations of the three pesticide formulations were cytotoxic and genotoxic in the *A. cepa* root tip meristem assay being reported. Because most adverse health effects by genotoxic agents are the result of genetic damage and any genetic activity of chemicals is most likely to result from cell division abnormalities, these genotoxic pesticides have potential to cause adverse environmental and health effects.

#### References

1. Aciole, E.H.P., Nilza, N. G., Andre, S. S., Erima, M. A., Sergio, M. N., Ana, C.L., Garcia, Kênya S. C. and Claudia, R. "Genetic toxicity of dillapiol and spinosad larvicides in somatic cells of *Drosophila melanogaster*." *Pest management science* 70.4 (2014): 559-565.
2. Akmoutsou, P., Despoina, M., Ifigeneia, N., Alexandros, O., Xrisa, P., Ilias, K., Gerasimos, F. et al. "Evaluation of toxicity and genotoxic effects of spinosad and deltamethrin in *Drosophila melanogaster* and *Bactrocera oleae*." *Pest management science* 67.12 (2011): 1534-1540.
3. Aktar, M.W., Dwaipayan, S. and Ashim, C. "Impact of pesticides use in agriculture: their benefits and hazards." *Interdisciplinary toxicology* 2.1 (2009): 1-12.
4. Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K. and Walter, P. "Molecular Biology of the Cell." *Garland Science, New York* 1601 (2008):1-1616.
5. Amer, S. M., and Odette, R. F. "Cytological effects of pesticides VI. Effect of the insecticide "Rogor" on the mitosis of *Vicia faba* and *Gossypium barbadense*." *Cytologia* 39.3 (1974): 507-514.
6. Aphicide Plus 100ml - Efekto. - <https://www.gardeningconcepts.co.za/products/aphicide-plus-100ml-efekto>
7. Asita, A.O., S. Moramang, T., Rants'o and S. Magama. "Modulation of mutagen-induced genotoxicity by vitamin C and medicinal plants in *Allium cepa* L." *Caryologia* 70.2 (2017): 151-165.
8. Aydemir, N., Serap, Ç., Şenay, S., Dilek, Y. and Özlem, Ö. "Evaluation of clastogenicity of 4, 6-dinitro-o-cresol (DNOC) in allium root tip test." (2008): 59-63.
9. Badr, A., A, Ghareeb. and H, M. El-Din. "Cytotoxicity of some pesticides in mitotic cells of *Vicia faba* roots." *Egypt. J. Appl. Sci* 7 (1992): 457-468.
10. Bianchi, J., Thais, C.C.F. and Maria, A. Marin-Morales. "Induction of mitotic and chromosomal abnormalities on *Allium cepa* cells by pesticides imidacloprid and sulfentrazone and the mixture of them." *Chemosphere* 144 (2016): 475-483.
11. Bonciu, E., Peter, F., Carmem, S. F., Jiang, W., Mehmet, C. K., Donghua, L., Felicia, M. et al. "An evaluation for the standardization of the *Allium cepa* test as cytotoxicity and genotoxicity assay." *Caryologia* 71.3 (2018): 191-209.
12. Bunch, T. R., C, Bond., K, Buhl. and D, Stone. "Spinosad General Fact Sheet; National Pesticide Information Center." *Oregon State University Extension Services* (2014).
13. Cai, D. W. "Understand the role of chemical pesticides and prevent misuses of pesticides." *Bulletin of Agricultural Science and Technology* 1.6 (2008): 36-38.
14. Chauhan, L. K. S., P, N. Saxena. and S, K. Gupta. "Cytogenetic effects of cypermethrin and fenvalerate on the root meristem cells of *Allium cepa*." *Environmental and experimental botany* 42.3 (1999): 181-189.
15. Chao, S.L., and John, E. C. "Interaction of imidacloprid metabolites and analogs with the nicotinic acetylcholine receptor of mouse brain in relation to toxicity." *Pesticide Biochemistry and Physiology* 58.1 (1997): 77-88.
16. Chaudhari, Renu. and Kamal, K.S. "Evaluation of Bifenthrin genotoxicity on *Channa*

- punctatus." *International Journal of Pure and Applied Zoology* 3.4 (2015): 297-301.
17. Cloyd, R.A., Cindy, L. G. and Stephen, R. K. "Greenhouse pesticide mixtures for control of silverleaf whitefly (Homoptera: Aleyrodidae) and twospotted spider mite (Acari: Tetranychidae)." *Journal of Entomological Science* 42.3 (2007): 375-382.
  18. Constantin, M.J. and Elizabeth, T. O. "Introduction and perspectives of plant genetic and cytogenetic assays a report of the US environmental protection agency Gene-Tox program." *Mutation Research/Reviews in Genetic Toxicology* 99.1 (1982): 1-12.
  19. Costa, C., V, Silvani, A, Melchini, S, Catania, J, J. Heffron, A, Trovato. and R, D. Pasquale. "Genotoxicity of imidacloprid in relation to metabolic activation and composition of the commercial product." *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 672.1 (2009): 40-44.
  20. Demisia, G., Dimitris, V., Marina, G. and Demetrios, P. M. "Assessment of the genotoxicity of imidacloprid and metalaxyl in cultured human lymphocytes and rat bone-marrow." *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 634.1-2 (2007): 32-39.
  21. Efekto - Eco Fruit Fly Bait gF 120 Insecticide - 100ml. -<https://www.takealot.com/efekto-eco-fruit-fly-bait-gf-120-insecticide-100ml/PLID41712349>
  22. Efekto - Rosecare 3 RTU - 750ml. -<https://www.hadeco.co.za/efekto/efekto-rosecare-3-rtu-750ml/>
  23. El-Ghamery, A. A., M, A. El-Kholy. and M, A. A. El-Yousser. "Evaluation of cytological effects of Zn<sup>2+</sup> in relation to germination and root growth of *Nigella sativa* L. and *Triticum aestivum* L." *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 537.1 (2003): 29-41.
  24. European Food Safety Authority (EFSA) Conclusion on the peer review of the pesticide risk assessment of the active substance myclobutanil. *EFSA Journal* 8.10 (2010): 1682-1765.
  25. Federation Register. "Imidacloprid: Pesticide Tolerances for Emergency Exemptions." *Federation Register* 70.196(2005): 59268-59276.
  26. Feng, S., Zhiming, K., Xinming, W., Pingan, P. and Eddy, Y. Z. "Assessing the genotoxicity of imidacloprid and RH-5849 in human peripheral blood lymphocytes in vitro with comet assay and cytogenetic tests." *Ecotoxicology and environmental safety* 61.2 (2005): 239-246.
  27. Feng, S., Zhiming, Kong, X. W., Lirong, Z. and Pingan, P. "Acute toxicity and genotoxicity of two novel pesticides on amphibian, *Rana N. Hallowell*." *Chemosphere* 56.5 (2004): 457-463.
  28. Feretti, D., Ilaria, Z., Claudia, Z., Elisabetta, C., M, Moretti. and Silvano, M. "Allium cepa chromosome aberration and micronucleus tests applied to study genotoxicity of extracts from pesticide-treated vegetables and grapes." *Food Additives and Contaminants* 24.6 (2007): 561-572.
  29. Gervais, J. A., et al. "Imidacloprid technical fact sheet. National Pesticide Information Center, Oregon State University Extension Services." (2010).
  30. Girotti, S., Elida, N.F. and Ermanno, E. "Persistence and distribution of pesticide residues in fresh agricultural food consumed in the province of Bologna." *Journal of Environmental Science and Health Part B* 44.2 (2009): 130-136.
  31. Graillot, V., Florence, T., Jean-Pierre, Cravedi. and Marc, A. "Evidence of the in vitro genotoxicity of methyl-pyrazole pesticides in human cells." *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 748.1-2 (2012): 8-16.
  32. Hayma, J. "The Storage of Agricultural Products (4th ed.)" *Agromisa Foundation. Netherlands* (2003).
  33. Hester, S., Tanya, M., William, T. P., Lynea, M., Charles, E. W. and Stephen, N. "The hepatocarcinogenic conazoles: cyproconazole, epoxiconazole, and propiconazole induce a common set of toxicological and transcriptional responses." *Toxicological Sciences* 127.1 (2012): 54-65.

34. Hester, S.D., Douglas, C.W., Stephen, N. and Sheau-Fung, T. "Transcriptional profiles in liver from rats treated with tumorigenic and non-tumorigenic triazole conazole fungicides: propiconazole, triadimefon, and myclobutanil." *Toxicologic pathology* 34.7 (2006): 879-894.
35. Higashi, K. "Metabolic activation of environmental chemicals by microsomal enzymes of higher plants." *Mutation Research* 19.2(1988): 273-288.
36. Hileman, B. "Environmental estrogens linked to reproductive abnormalities and cancer." *Chemical and Engineering News* 72.5 (1994): 19-23.
37. Hoppin, Jane A., et al. "Pesticides and atopic and nonatopic asthma among farm women in the Agricultural Health Study." *American journal of respiratory and critical care medicine* 177.1 (2008): 11-18.
38. "How does myclobutinil work?" Retrieved from [http://www.domyown.com/myclobutinil-c-114\\_468.html](http://www.domyown.com/myclobutinil-c-114_468.html). Iganci, J. R. V., V. L. Bobrowski., G. Heiden., V. C. Stein. and B. H. G. Rocha. "Efeito do extrato aquoso de diferentes espécies de boldo sobre a germinação e índice mitótico de *Allium cepa* L." *Arq Inst Biol* 73.1 (2006): 79-82.
39. Ilyushina, N.A., Olga, V. E., Gleb, V. M., Nataliya, S. A., Yulia, A. R., Valerii, N. R., Marina, G., Alexander V., Polychronis, S. and Aristidis, T. "Genotoxicity of mixture of imidacloprid, imazalil and tebuconazole." *Toxicology Reports* 7 (2020): 1090-1094.
40. Jemec, Anita., et al. "Comparative toxicity of imidacloprid, of its commercial liquid formulation and of diazinon to a non-target arthropod, the microcrustacean *Daphnia magna*." *Chemosphere* 68.8 (2007): 1408-1418.
41. Jeyaratnam, J. "Acute pesticide poisoning: a major global health problem." *World Health Statistics Quarterly* 43.3 (1990): 139-144.
42. Jha, R.K. and A.P. Regmi. "Productivity of pesticides in vegetable farming in Nepal." *The South Asian Network for Development and Environmental Economics (SANDEE)* (2009): 42-09.
43. Johnson, M., B, Luukinen., J, Gervais., K, Buhl. and D, Stone. "Bifenthrin Technical Fact Sheet. National Pesticide Information Center." *Oregon State :University Extension Services.* (2010).
44. Kamel, F., L, S. Engel., B, C. Gladen., J, A. Hoppin., M, C.R. Alavanja. and D, P. Sandler. "Neurologic symptoms in licensed pesticide applicators in the Agricultural Health Study." *Human & experimental toxicology* 26.3 (2007): 243-250.
45. Karabay, N. U., and M, G.O. "Cytogenetic and genotoxic effects of the insecticides, imidacloprid and methamidophos." *Genetics and Molecular Research* 4.4 (2005): 653-662.
46. Kreutzweiser, D., Kevin, G., Derek, C., Taylor, S. and Dean, T. "Non-target effects on aquatic decomposer organisms of imidacloprid as a systemic insecticide to control emerald ash borer in riparian trees." *Ecotoxicology and Environmental Safety* 68.3 (2007): 315-325.
47. Leme, D.M. and Maria, A., Marin-Morales. "Allium cepa test in environmental monitoring: a review on its application." *Mutation Research/Reviews in Mutation Research* 682.1 (2009): 71-81.
48. Levan, A. "The effect of colchicine on root mitoses in *Allium*." *Hereditas* 2.4(1938): 471-486.
49. Lewis, K.A., et al. "An international database for pesticide risk assessments and management." *Human and Ecological Risk Assessment: An International Journal* 22.4 (2016): 1050-1064.
50. Linnainmaa, K., et al. "Cytogenetic effects of styrene and styrene oxide." *Mutation Research/Genetic Toxicology* 58.2-3 (1978): 277-286.
51. Liu, C. J., et al. "The pollution of pesticides in soils and its bioremediation." *System Sciences and Comprehensive Studies in Agriculture* 18.4 (2002): 295-297.
52. Mansour, S. A., A. H. Mossa. and T, M. Heikal. "Cytogenetic and hormonal alteration in rats exposed to recommended "safe doses" of spinosad and malathion insecticides." *Int J Agric Biol* 10.1 (2008): 9-14.
53. Mansour, S. "Pesticide exposure-Egyptian scene." *Toxicology* 198 1-3(2004): 91-115.

54. Marcano, L., et al. "Cytotoxicity and mode of action of maleic hydrazide in root tips of *Allium cepa* L." *Environmental Research* 94.2 (2004): 221-226.
55. Marrs, T. C. (Ed.) "Mammalian Toxicology of Insecticides." *Royal Society of Chemistry* (2012): 1-49.
56. Medina, D., et al. "Persistence of organophosphorus pesticide residues in Limon River waters." *Bulletin of Environmental Contamination and Toxicology* 63.1 (1999): 39-44.
57. Mendonça, T.P., Jéssica, D.de. A., Weverson, J. S., Daniele, R.M., Carlos, F.C., Jéssica, S.V., Nathalya P.B., et al. "Genotoxic and mutagenic assessment of spinosad using bioassays with *Tradescantia pallida* and *Drosophila melanogaster*." *Chemosphere* 222 (2019): 503-510.
58. Mertz, F.P. and R.C. Yao. "Saccharopolyspora spinosa sp. nov. isolated from soil collected in a sugar mill rum still." *International Journal of Systematic and Evolutionary Microbiology* 40.1 (1990): 34-39.
59. Norppa, H. "Cytogenetic biomarkers and genetic polymorphisms." *Toxicology letters* 149.1-3 (2004): 309-334.
60. Österberg, R., Dan, P. and Gunnar, B. "The condensation of DNA by chromium (III) ions." *Journal of Biomolecular Structure and Dynamics* 2.2 (1984): 285-290.
61. Panda, B. B., and U, K. Sahu. "Induction of abnormal spindle function and cytokinesis inhibition in mitotic cells of *Allium cepa* by the organophosphorus insecticide fensulfothion." *Cytobios* 42.167-168 (1985): 147-155.
62. Parry, E. M., L, E. Mumba., A, Asita. And J, M. Parry. "Mechanisms of Action of Aneuploidy Inducing Chemicals with Particular Reference to Spindle Inhibitors and Neurotoxins." In R. C. Sobti, G. Obe & P. Quillardet (Eds), "Trends in environmental mutagenesis." *Tausco Book distributors, New Delhi, India, Publisher* (1999): 101-110
63. Parry, E. M., D, C. Sharp. and James, M. P. "The observation of mitotic division aberrations in mammalian cells exposed to chemical and radiation treatments." *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis* 150.1-2 (1985): 369-381.
64. GI, Bhat. "A comparative study of MH and EMS in the induction of chromosomal aberrations on lateral root meristem in *Clitoria ternatea* L." *Cytologia* 57.2 (1992): 259-264.
65. Pimentel, D. H., Acquay, M. B., P, Rice. M., Silva, J. N., V, Lipner., S, Giordano., A. Horowitz. and M, D'amore. "Assessment of environmental and economic impacts of pesticide use." *The pesticide question*. Springer, Boston, MA, 1993. 47-84.
66. Plewa, M. J., and James, M. G. "The activation of chemicals into mutagens by green plants." *Chemical Mutagens*. Springer, Boston, MA, 1982. 401-420.
67. Raja, N., et al. "Potential of plants for protecting stored pulses from *Callosobruchus maculatus* (Coleoptera: Bruchidae) infestation." *Biological agriculture & horticulture* 19.1 (2001): 19-27.
68. Reddy, S.E.G., N,T. Rao. and K,S. Devi. "Genotoxicity evaluation of bifenthrin pyrethroid insecticide to freshwater fish *Danio rerio* using the alkaline Comet Assay in gill tissue." *Indian Journal of Applied Research* 3.12 (2013): 53-56.
69. Richter, E. D. "Acute human pesticide poisonings." *Encyclopedia of pest management* 1 (2002): 3-6.
70. Robson, M. G. "Global Pesticide Use: Weighing the risk and benefits." *World Ecology Report, Spring, World Information Transfer, New York* 29.1 (2019): 1-16.
71. Rodríguez, Y.A., Cintya, A.C., Janaína, P., Odair, C.B., Osmar, M., Rafael, A.C.F. and Carmem, S. F. "*Allium cepa* and *Tradescantia pallida* bioassays to evaluate effects of the insecticide imidacloprid." *Chemosphere* 120 (2015): 438-442.
72. Rojas, A., María, E. O. and Ximena, B. "Congenital malformations and pesticide exposure." *Revista médica de Chile* 128.4 (2000): 399-404.
73. Ross, J.A., Tanya, M. and Sharon, A. L. "In vivo mutagenicity of conazole fungicides cor-

- relates with tumorigenicity." *Mutagenesis* 24.2 (2009): 149-152.
74. Ross, J.A., Sharon, A. L., Judith, E. S. and Garret, B. N. "Quantitative changes in endogenous DNA adducts correlate with conazole in vivo mutagenicity and tumorigenicity." *Mutagenesis* 27.5 (2012): 541-549.
75. Schulz-Jander., Daniel, A. and John, E. C. "Imidacloprid insecticide metabolism: human cytochrome P450 isozymes differ in selectivity for imidazolidine oxidation versus nitroimine reduction." *Toxicology letters* 132.1 (2002): 65-70.
76. Shah, R.G., Jean, L., Sanjay, K., Patrick, L., Pierre, A., Martine, T., John, Z. and Guy, G. Poirier. "Determination of genotoxicity of the metabolites of the pesticides Guthion, Sencor, Lorox, Reglone, Daconil and Admire by 32P-postlabeling." *Molecular and cellular biochemistry* 169.1 (1997): 177-184.
77. Sheets, L. P. "Imidacloprid: A Neonicotinid Insecticide." *Handbook of Pesticide Toxicology*, 2nd ed.; Krieger, R. I., Ed.; Academic Press: San Diego, CA 2.54 (2001): 1123-1130.
78. Sheets, L. P. "The Neonicotinoid Insecticides." *Handbook of Neurotoxicology*; Massaro, E. J., Ed.; Humana Press: Totowa, NJ 1 (2002); 79-87.
79. Smaka-Kincl, V., Peter, Stegnar, M.L. and Mihael, J. T. "The evaluation of waste, surface and ground water quality using the Allium test procedure." *Mutation Research/Genetic Toxicology* 368.3-4 (1996): 171-179.
80. Solecki, R. "Toxicological evaluations. Imidacloprid." *Joint Meeting on Pesticide Residues*. 2001.
81. Stebbins, K. E. "Spinosad insecticide: subchronic and chronic toxicity and lack of carcinogenicity in CD-1 mice." *Toxicological Sciences*, 65.2(2002): 276-287.
82. Subbarao, N. S. "Soil Microbiology (4th ed.)" *Science Publishers, Inc* (1999): 303-324.
83. Sultana, S., Asma, A.P.S., Kaiser, J. and Abbas, H. A. "Evaluation of cytotoxicity and genotoxicity of pesticide mixtures on lymphocytes." *Toxicology mechanisms and methods* 26.8 (2016): 588-594.
84. Tedesco, S.B. and Haywood, L IV. "Bioindicator of genotoxicity: The Allium cepa test." *Environmental contamination* (2012): 137-156.
85. Thyssen, J. and L, Macheimer. "Imidacloprid: Toxicology and Metabolism." In Yamamoto, I. & Casida, J.E. (eds). "Nicotinoid Insecticides and the Nicotinic Acetylcholine Receptor." *Springer, Tokyo* (1999): 213-222.
86. Tomlin, C. D. S. "The Pesticide Manual, A World Compendiumpp." 12th ed. *British Crop Protection Council: Surry, England* (2000):502-504.
87. Trapp, Kassia Cauana, C. A. L. Hister, H. D. Laughinghouse IV, Aline Augusti Boligon, and Solange Bosio Tedesco. "Determination of phenolic compounds and evaluation of cytotoxicity in *Plectranthus barbatus* using the Allium cepa test." *Caryologia* 73.2 (2020): 145-153.
88. U.S Environmental Protection Agency (EPA) "Bifenthrin Pesticide Fact Sheet" *Office of Prevention, Pesticides, and Toxic Substances, Office of Pesticide Programs, U.S. Government Printing Office: Washington, DC* (1988).
89. U.S Environmental Protection Agency (EPA) "Spinosad Pesticide Fact Sheet No. HJ 501C." *Office of Pesticides and Toxic Substances, Washington, DC* (1997).
90. U.S Environmental Protection Agency (EPA) "Myclobutanil: Pesticide Tolerances." 65 FR 29963, *Office of Pesticide Programs, U.S. Government Printing Office, Washington, DC* (2000).
91. Vieira, A. F. C. and Graciele, L.S. "Cyto (geno) toxic endpoints assessed via cell cycle bioassays in plant models." *Cytotoxicity. London: Intech Open* (2018): 117-129.
92. WHO- World Health Organisation Report. "WHO Specifications and Evaluations for Public Health Pesticides." (2005).
93. Spinosad: Evaluation Report 636(2005): 1-20.
94. WHO-World Health Organization Report. "WHO Specifications and Evaluations for Public Health Pesticides: Bifenthrin." 2017.
95. Yano, B. L., D, M. Bond., M, N. Novilla., L, G. McFadden. and M, J. Reasor. "Spinosad insecticide: subchronic and chronic toxicity and

- lack of carcinogenicity in Fischer 344 rats." *Toxicological Sciences* 65.2 (2002): 288-298.
96. Mustafa, Y. and Evrim, S.A. "Genotoxicity testing of quizalofop-P-ethyl herbicide using the *Allium cepa* anaphase-telophase chromosome aberration assay." *Caryologia* 61.1 (2008): 45-52.
97. Zang, Y., Y, Zhong., Y, Luo. and Z, M. Kong. "Genotoxicity of two novel pesticides for the earthworm, *Eisenia fetida*." *Environmental Pollution* 108.2 (2000): 271-278.
98. Zhang, A., Hartmut, K., Peter, Maienfisch. and John, E. C. "Insect Nicotinic Acetylcholine Receptor: Conserved Neonicotinoid Specificity of [3H] Imidacloprid Binding Site: Conserved Neonicotinoid Specificity of [3H] Imidacloprid Binding Site." *Journal of neurochemistry* 75.3 (2000): 1294-1303.
99. Zhang, C.Z., Alexander, S., Hauke, C., Joshua, M.F., Emily, K. J., Shiwei, L., Matthew, M. and David, P. "Chromothripsis from DNA damage in micronuclei." *Nature* 522.7555 (2015): 179-184.

**Source of support:** Nil; **Conflict of interest:** Nil.

**Cite this article as:**

Asita, O. A., Relebohile, R. M. and Sibusisiwe, M. "Genotoxicity of Aphicide Plus®, Eco Fruit-fly Bait GF120® and Rosecare 3® in the *Allium cepa* assay." *Annals of plant sciences* 10.12 (2021):pp 4436-4449